

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/280740830>

Breeding bird community of the Montagne della Duchessa (Latium, central Italy)

Article in *Avocetta* - June 2015

CITATIONS

3

READS

160

5 authors, including:



Massimo Brunelli

SROPU

70 PUBLICATIONS 246 CITATIONS

[SEE PROFILE](#)



Michele Cento

Stazione Romana Osservazione e Protezione Uccelli (SROPU)

59 PUBLICATIONS 81 CITATIONS

[SEE PROFILE](#)



Emiliano De Santis

Monti Simbruini Natural Regional Park - Lazio

14 PUBLICATIONS 24 CITATIONS

[SEE PROFILE](#)



Enzo Savo

13 PUBLICATIONS 18 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Italy *Ciconia nigra* monitoring [View project](#)



Radar monitoring and visual observations at the Strait of Messina [View project](#)

Breeding bird community of the Montagne della Duchessa (Latium, central Italy)

MASSIMO BRUNELLI^{1*}, MICHELE CENTO¹, EMILIANO DE SANTIS², ENZO SAVO¹, ALBERTO SORACE¹

¹Stazione Romana Osservazione e Protezione Uccelli, Roma, Italy

²Servizio Monitoraggio e Sorveglianza, Parco Nat. Reg. Monti Simbruini, Jenne (Roma), Italy

*Corresponding author: mss.brunelli@tin.it

Abstract – During 2014 spring we studied the breeding bird community of ‘Montagne della Duchessa’ Regional Nature Reserve (Latium, Italy). In the study area (5870 ha; altitude range: 800-2239 m a.s.l.) we performed 127 point counts (10 min long) among four macro-habitat types (arable crop, oak wood, beech wood, bushes and alpine meadow) distributed proportionally to their respective area. We found 88 breeding species (79 detected during point counts). Most abundant species were: *Erithacus rubecula*, *Sylvia atricapilla* and *Corvus cornix*. Most widespread species were *Erithacus rubecula*, *Fringilla coelebs*, *Sylvia atricapilla*, *Turdus merula*, *Phylloscopus collybita* and *Cyanistes caeruleus*. In arable crops we detected a total of 54 species (*Passer italiae*, *Sturnus vulgaris*, *Corvus cornix* and *Passer montanus* the most abundant); in oak woods 34 species (*Erithacus rubecula*, *Turdus merula*, *Cyanistes caeruleus*, *Sylvia atricapilla*, *Parus major*, *Phylloscopus collybita*, *Poecile palustris* and *Fringilla coelebs* the most abundant); in beech woods 37 species (*Erithacus rubecula*, *Fringilla coelebs*, *Sylvia atricapilla*, *Poecile palustris*, *Cyanistes caeruleus*, *Sitta europaea* and *Phylloscopus collybita* the most abundant); in bushes and alpine meadows 33 species (*Oenanthe oenanthe*, *Anthus spinoletta*, *Alauda arvensis*, *Carduelis cannabina* and *Anthus campestris* the most abundant). In bushes and alpine meadows Richness, Abundance and Diversity were lowest. The collected data showed the important role of this Protected Area for some species of conservation concern.

Key-words: birds community, richness, abundance, dominance, diversity, evenness, central Italy.

INTRODUCTION

The present contribution was carried out for a wider project aimed at the realization of the Atlas of breeding birds in the Montagne della Duchessa Regional Nature Reserve and surrounding areas. Bird atlases are an useful tool to obtain data on distribution and local ecology of the species (e.g. Sutherland *et al.* 2004, Devictor & Robert 2009, Sorace & Gustin 2010, Belmaker *et al.* 2012). Therefore, several regional protected areas have accomplished or undertaken similar projects (Taffon *et al.* 2008, De Santis *et al.* 2009, Ukmar *et al.* 2009, Papi *et al.* 2010, Battisti & Guidi 2012, Quatrini *et al.* 2012).

The management of a natural protected area needs basic cognitive tools that on one side can help to address the planning of the activities and on the other side can constitute the elements of successive monitoring to assess the effectiveness of the conservation and management actions that were previously undertaken (Primack & Carotenuto 2003, Apollonio *et al.* 2004).

Besides its role as management tool, this work, represents also an important contribution to the knowledge of

the breeding avifauna in this area of central Apennines. Excluding the information at regional scale of the recent Atlas of Latium breeding birds (Brunelli *et al.* 2011), studies on local birds are scarce, often dated or focused on single bird group or species (Rossi 1947, Di Carlo 1980, Spinetti 1997, Allavena *et al.* 1999, Allavena & Panella 2000, D.R.E.A.M. Italia 2003, AA.VV. 2004, Bernoni *et al.* 2009, Sorace *et al.* 2011). In addition, few investigations deal with composition and structure of breeding bird communities of massifs and mountain chains of central and southern Italy, often referring to a single environment (De Filippo & Kalby 1985, De Filippo *et al.* 1985, Mingozzi 1994, Brandmayr *et al.* 1996, Vuerich *et al.* 2006, Fulco & Tellini Florenzano 2008, Mancinelli *et al.* 2009, Battisti *et al.* 2010, Sorace 2011, Strinella *et al.* 2011).

STUDY AREA

The study area, placed in the Borgorose district (RI) in the eastern Latium at the border of Abruzzo region (Fig. 1), extends over about 5870 ha including 3547 ha inside




Figure 1. Study area with the overlapped 1-km square grid. In grey, the Regional Nature Reserve “Montagne della Duchessa”.

the Montagne della Duchessa Regional Nature Reserve and 2323 ha in the adjoining areas with higher naturalistic value, in particular the Corvaro plain and the slopes of Cava Mountain towards Malito Valley. Altitude ranges between 800 m a.s.l. of Corvaro plain and 2239 m of Costone Mountain.

Due to the presence of species and habitat of European Community interest, Montagne della Duchessa district includes three sites of Natura 2000 network: two SCI, instituted according to Habitat Directive (92/43/CEE), named “Monti della Duchessa (area sommitale)” (IT6020020) and “Monte Duchessa-Vallone Cieco e Bosco Cartore” (IT 6020021); the two SCI are included in ZSP “Riserva Naturale Montagne della Duchessa” (IT3020046), established according to Birds Directive (2009/147/CEE) and overlapping with the Nature Reserve that was instituted by Latium Regional Law in 1990. The nearby areas of Cava Mountain, Malito Valley and Corvaro plain are not protected.

At the lower altitude (800-1100 m a.s.l.) the vegetation of study area is characterized by the presence of farmland and meadows/pastures, in particular in Corvaro plain and, partially, in Ruara Valley and Malito Valley, and by large areas covered by bushes (mainly with *Rosa canina*, *Prunus spinosa*, *Cornus mas*) and oak woods (with *Quercus cerris* and *Quercus pubescens* dominants). Moreover, wide chestnut groves with *Castanea sativa* are present in Malito Valley while *Salix alba* e *Populus nigra* are common along Apa stream. At the higher altitude (1200-1900 m) oak woods are gradually replaced by mixed woods with *Fagus sylvatica*. Pure beech woods prevail between 1500 and 1800 m and above them extensive bushes with *Juniperus communis* alternate with alpine xeric meadows

(for a more detailed description of the study area see Russo *et al.* 2010).

METHODS

In order to make the investigation replicable for future monitoring and to obtain representative data, we used a standardized methodology widely experimented in analogous ornithological investigations (see Fornasari *et al.* 2010). We overlapped a 1-km square grid, originated from UTM coordinate system, on the map of the study area obtaining 63 recording units (RU) identified by an univocal alpha-numeric denomination (Fig. 1).

We investigated the structure of breeding bird communities by means of point count method (Blondel *et al.* 1970, Bibby *et al.* 2000, Fornasari *et al.* 2010). Individuals of more rare or less detectable species might be not sampled with this method, nevertheless collected data with point counts can well describe bird communities, characterizing their variation throughout the environmental mosaic (Sutherland *et al.* 2004, Battisti *et al.* 2010).

We carried out 127 point counts, not less than two per each of the 63 RU, spaced at least 200 m apart. Point counts were located by means of stratifying sampling using the Software Quantum Gis 2.2 and the CORINE Land Cover categories (Regione Lazio 2010). We selected four main habitat types:

- arable crops (23 point counts);
- oak wood (22 point counts);
- beech wood (46 point counts);
- bushes and alpine meadow (36 point counts).

Some scarcely represented habitats, such as urban areas (continuous and discontinuous urban fabric, road network, etc) and chestnut groves, were incorporated with the main four habitats. When a point count was not achievable, due to the uneven morphology of the study area, it was replaced by a new point in the same habitat and RU.

In 2014 breeding season, each point count (10-min long), was double visited recording the species of all birds identifiable by sight or song inside and outside a 100-m radius from the observer. The first visit was carried out between 15 March and 18 May, the second one between 20 May and 1 July; time lag between two visits in each point count was at least 35 days. One point was assigned to each recorded individual. For each species, the maximum score obtained between the two visits was considered the number of individuals present per point count.

To characterize the structure of breeding bird communities, we calculated the following parameters:

- Richness (S): total number of recorded species;
- Richness mean (Smean): mean number of recorded species per point count;
- Abundance (A): mean number of individuals per point count;
- Dominance (p_i): where p_i is the relative frequency of each species; species with $p_i \geq 0.050$ and $p_i \geq 0.020$ were considered respectively dominant and subdominant (Turcek 1956);
- Diversity (H): $H = -\sum p_i \ln p_i$ (Shannon & Weaver 1963);
- Evenness (J): $J = H/\ln S$ (Lloyd & Ghelardi 1964, Pielou 1966);
- Non-Passeriformes percentage (Ferry & Frochot 1970).

To analyze data concerning the community parameters and the frequencies of species in the whole study area, we summed data collected in each point count both inside and outside the 100-m radius from the observer. In the analysis in which bird communities of different habitats were separately considered, only data collected inside the 100-m radius were taken into account.

Since data considered in statistical analysis were not normally distributed (Kolmogorov-Smirnov test), we used non-parametric tests: the non-Passeriformes percentages were compared with χ^2 test; the differences of Richness e Abundance values per point count among the four habitats were evaluated by means of Kruskal-Wallis ANOVA rank Test with Mann-Whitney U Test for pairwise comparisons; the differences of Dominance of recorded species among the four habitats were tested by means of Friedman test with Wilcoxon test for pairwise comparisons.

Besides collected data with point counts, we carried out some not standardized field researches to obtain infor-

mation about less detectable breeding species. We did not take into account the information collected for these species in the computation of the community parameters.

RESULTS

On the whole we recorded 88 breeding species, including 23 non Passerines (26.1%) and 65 Passerines (73.9%); 79 out of 88 species were observed with the 127 point counts, including 20 non Passerines (25.3%) and 59 Passerines (74.7%) (Tab. 1). The dominant species resulted in decreasing order *Erithacus rubecula*, *Sylvia atricapilla* and *Corvus cornix*; 17 were the subdominant species (Tab. 1). The most widespread species in the study area, i.e. recorded at least in 50% of point counts, resulted in decreasing order *Erithacus rubecula*, *Fringilla coelebs*, *Sylvia atricapilla*, *Turdus merula*, *Phylloscopus collybita* and *Cyanistes caeruleus* (Tab. 1).

In Table 2 the values of breeding bird community parameters of the study area are shown.

In Table 3, data on Abundance and Dominance of the 73 species recorded inside the 100-m radius in the four habitat types are reported (see Methods).

In the arable crops we observed 54 species, among them *Passer italiae*, *Sturnus vulgaris*, *Corvus cornix* and *Passer montanus*, species typical of man-made environments, resulted dominant. Other ten species were subdominant, including mainly ecotonal species, but also *Pica pica*, *Galerida cristata* and *Emberiza calandra*, which are linked to agricultural environments.

In the oak wood we found 34 species, among them *Erithacus rubecula*, *Turdus merula*, *Cyanistes caeruleus*, *Sylvia atricapilla*, *Parus major*, *Phylloscopus collybita*, *Poecile palustris* and *Fringilla coelebs* resulted dominant in decreasing order. These eight species are typical of not particularly old forests. Species exclusive of this habitat were not observed.

In the beech wood we recorded 37 species, among them *Erithacus rubecula*, *Fringilla coelebs*, *Sylvia atricapilla*, *Poecile palustris*, *Cyanistes caeruleus*, *Sitta europaea* and *Phylloscopus collybita* resulted dominant. Four species were only found in this habitat: *Emberiza cia* (in edge zone), *Phylloscopus sibilatrix*, *Certhia familiaris* and *Ficedula albicollis*. The recording of the last three species and the observations of three Picidae species indicate the availability of mature wood sectors.

In bushes and alpine meadows we found 33 species, among them *Oenanthe oenanthe*, *Anthus spinoletta*, *Alauda arvensis*, *Carduelis cannabina* and *Anthus campestris*, typical of open habitats, were dominant in decreas-

Table 1. Number of point counts in which each species has been detected and mean number of individuals per point count (\pm SD) in spring 2014. In bold, dominant species ($\pi > 0.05$); in italic, sub-dominant species ($\pi > 0.02$). Threat categories are evaluated on the basis of Bird Directive Annex I (2009/147/CEE), SPEC classification (BirdLife International 2004), National Red List (Peronace *et al.* 2012) and Regional Red List (Calvario *et al.* 2011) categories. Specie marked with * were found only outside point counts.

| Species | Occupied points | | Ind./point counts | | | Bird Directive | SPEC | National Red List | Regional Red List |
|-----------------------------------|-----------------|-------|-------------------|-------|--------------|----------------|-----------------------|-------------------|-------------------|
| | N | % | Mean | SD | π | | | | |
| <i>Anas platyrhynchos</i> | 1 | 0.79 | 0.024 | 0.266 | 0.001 | | Non-SPEC | LC | |
| <i>Alectoris graeca</i> * | - | - | - | - | - | I | SPEC2 | VU | VU |
| <i>Coturnix coturnix</i> | 2 | 1.57 | 0.016 | 0.125 | 0.001 | II B | SPEC3 | DD | |
| <i>Pernis apivorus</i> | 2 | 1.57 | 0.016 | 0.125 | 0.001 | I | Non-SPEC ^E | LC | VU |
| <i>Gyps fulvus</i> | 13 | 10.24 | 0.417 | 1.725 | 0.018 | I | Non-SPEC | CR | CR |
| <i>Circaetus gallicus</i> * | - | - | - | - | - | I | SPEC3 | VU | EN |
| <i>Accipiter nisus</i> | 2 | 1.57 | 0.016 | 0.125 | 0.001 | | Non-SPEC | LC | |
| <i>Buteo buteo</i> | 15 | 11.81 | 0.181 | 0.583 | 0.008 | | Non-SPEC | LC | |
| <i>Falco tinnunculus</i> | 12 | 9.45 | 0.094 | 0.294 | 0.004 | | SPEC3 | LC | |
| <i>Falco peregrinus</i> | 3 | 2.36 | 0.024 | 0.152 | 0.001 | I | Non-SPEC | LC | NT |
| <i>Columba livia</i> f. domestica | 2 | 1.57 | 0.016 | 0.125 | 0.001 | | Non-SPEC | - | |
| <i>Columba palumbus</i> | 55 | 43.31 | 0.606 | 0.828 | 0.026 | II A | Non-SPEC ^E | LC | |
| <i>Streptopelia decaocto</i> | 1 | 0.79 | 0.016 | 0.177 | 0.001 | | Non-SPEC | LC | |
| <i>Streptopelia turtur</i> | 5 | 3.94 | 0.039 | 0.195 | 0.002 | II B | SPEC3 | LC | |
| <i>Cuculus canorus</i> | 41 | 32.28 | 0.378 | 0.603 | 0.016 | | Non-SPEC | LC | |
| <i>Strix aluco</i> | 2 | 1.57 | 0.016 | 0.125 | 0.001 | | Non-SPEC ^E | LC | |
| <i>Caprimulgus europaeus</i> * | - | - | - | - | - | I | SPEC2 | LC | DD |
| <i>Apus apus</i> | 1 | 0.79 | 0.039 | 0.444 | 0.002 | | Non-SPEC | LC | |
| <i>Upupa epops</i> | 7 | 5.51 | 0.063 | 0.275 | 0.003 | | SPEC3 | LC | |
| <i>Jynx torquilla</i> | 3 | 2.36 | 0.024 | 0.152 | 0.001 | | SPEC3 | EN | |
| <i>Picus viridis</i> | 37 | 29.13 | 0.339 | 0.567 | 0.015 | | SPEC2 | LC | |
| <i>Dendrocopos major</i> | 17 | 13.39 | 0.142 | 0.372 | 0.006 | | Non-SPEC | LC | |
| <i>Dendrocopos minor</i> | 2 | 1.57 | 0.016 | 0.125 | 0.001 | | Non-SPEC | LC | DD |
| <i>Galerida cristata</i> | 8 | 6.30 | 0.126 | 0.563 | 0.005 | | SPEC3 | LC | |
| <i>Lullula arborea</i> | 10 | 7.87 | 0.110 | 0.422 | 0.005 | I | SPEC2 | LC | |
| <i>Alauda arvensis</i> | 32 | 25.20 | 0.866 | 3.105 | 0.037 | II B | SPEC3 | VU | |
| <i>Ptyonoprogne rupestris</i> * | - | - | - | - | - | | Non-SPEC | LC | |
| <i>Hirundo rustica</i> | 3 | 2.36 | 0.079 | 0.543 | 0.003 | | SPEC3 | NT | |
| <i>Delichon urbicum</i> | 6 | 4.72 | 0.205 | 1.197 | 0.009 | | SPEC3 | NT | |
| <i>Anthus campestris</i> | 16 | 12.60 | 0.228 | 0.657 | 0.010 | I | SPEC3 | LC | DD |
| <i>Anthus trivialis</i> | 28 | 22.05 | 0.346 | 0.770 | 0.015 | | Non-SPEC | VU | |
| <i>Anthus spinoletta</i> | 28 | 22.05 | 0.496 | 1.076 | 0.021 | | Non-SPEC | LC | |
| <i>Motacilla cinerea</i> | 1 | 0.79 | 0.008 | 0.089 | 0 | | Non-SPEC | LC | |
| <i>Motacilla alba</i> | 6 | 4.72 | 0.063 | 0.327 | 0.003 | | Non-SPEC | LC | |
| <i>Troglodytes troglodytes</i> | 45 | 35.43 | 0.480 | 0.754 | 0.021 | | Non-SPEC | LC | |
| <i>Prunella collaris</i> * | - | - | - | - | - | | Non-SPEC | LC | DD |
| <i>Erithacus rubecula</i> | 84 | 66.14 | 1.646 | 1.551 | 0.071 | | Non-SPEC ^E | LC | |
| <i>Luscinia megarhynchos</i> | 22 | 17.32 | 0.260 | 0.693 | 0.011 | | Non-SPEC ^E | LC | |
| <i>Phoenicurus ochruros</i> | 13 | 10.24 | 0.142 | 0.483 | 0.006 | | Non-SPEC | LC | |
| <i>Phoenicurus phoenicurus</i> | 9 | 7.09 | 0.110 | 0.422 | 0.005 | | SPEC2 | LC | VU |
| <i>Saxicola torquatus</i> | 7 | 5.51 | 0.071 | 0.313 | 0.003 | | Non-SPEC | VU | |
| <i>Oenanthe oenanthe</i> | 29 | 22.83 | 0.567 | 1.288 | 0.024 | | SPEC3 | NT | |
| <i>Monticola saxatilis</i> | 5 | 3.94 | 0.047 | 0.247 | 0.002 | | SPEC3 | VU | EN |

continued

Breeding bird community of the Montagne della Duchessa

| Species | Occupied points | | Ind./point counts | | | Bird Directive | SPEC | National Red List | Regional Red List |
|--------------------------------------|-----------------|-------|-------------------|-------|--------------|----------------|-----------------------|-------------------|-------------------|
| | N | % | Mean | SD | pi | | | | |
| <i>Turdus merula</i> | 71 | 55.91 | 0.882 | 1.081 | 0.038 | II B | Non-SPEC ^E | LC | |
| <i>Turdus philomelos</i> | 9 | 7.09 | 0.094 | 0.366 | 0.004 | II B | Non-SPEC ^E | LC | |
| <i>Turdus viscivorus</i> | 27 | 21.26 | 0.260 | 0.580 | 0.011 | II B | Non-SPEC ^E | LC | |
| <i>Sylvia atricapilla</i> | 80 | 62.99 | 1.339 | 1.387 | 0.058 | | Non-SPEC ^E | LC | |
| <i>Sylvia communis</i> * | - | - | - | - | - | | Non-SPEC ^E | LC | |
| <i>Sylvia cantillans</i> | 4 | 3.15 | 0.039 | 0.232 | 0.002 | | Non-SPEC ^E | LC | |
| <i>Phylloscopus bonelli</i> | 27 | 21.26 | 0.346 | 0.749 | 0.015 | | SPEC2 | LC | |
| <i>Phylloscopus sibilatrix</i> | 4 | 3.15 | 0.055 | 0.341 | 0.002 | | SPEC2 | LC | |
| <i>Phylloscopus collybita</i> | 69 | 54.33 | 0.921 | 1.066 | 0.040 | | Non-SPEC | LC | |
| <i>Regulus ignicapilla</i> | 6 | 4.72 | 0.055 | 0.261 | 0.002 | | Non-SPEC ^E | LC | |
| <i>Muscicapa striata</i> * | - | - | - | - | - | | SPEC3 | LC | |
| <i>Ficedula albicollis</i> | 2 | 1.57 | 0.024 | 0.198 | 0.001 | I | Non-SPEC ^E | LC | |
| <i>Aegithalos caudatus</i> | 13 | 10.24 | 0.189 | 0.880 | 0.008 | | Non-SPEC | LC | |
| <i>Cyanistes caeruleus</i> | 65 | 51.18 | 0.787 | 1.005 | 0.034 | | Non-SPEC ^E | LC | |
| <i>Parus major</i> | 62 | 48.82 | 0.795 | 1.079 | 0.034 | | Non-SPEC | LC | |
| <i>Periparus ater</i> | 26 | 20.47 | 0.339 | 0.857 | 0.015 | | Non-SPEC | LC | |
| <i>Poecile palustris</i> | 41 | 32.28 | 0.512 | 0.872 | 0.022 | | SPEC3 | LC | |
| <i>Sitta europaea</i> | 42 | 33.07 | 0.567 | 0.964 | 0.024 | | Non-SPEC | LC | |
| <i>Tichodroma muraria</i> * | - | - | - | - | - | | Non-SPEC | LC | EN |
| <i>Certhia familiaris</i> | 4 | 3.15 | 0.039 | 0.232 | 0.002 | | Non-SPEC | LC | NT |
| <i>Certhia brachydactyla</i> | 7 | 5.51 | 0.071 | 0.313 | 0.003 | | Non-SPEC ^E | LC | |
| <i>Oriolus oriolus</i> | 5 | 3.94 | 0.047 | 0.247 | 0.002 | | Non-SPEC | LC | |
| <i>Lanius collurio</i> | 5 | 3.94 | 0.063 | 0.351 | 0.003 | I | SPEC3 | VU | |
| <i>Garrulus glandarius</i> | 42 | 33.07 | 0.528 | 0.958 | 0.023 | II B | Non-SPEC | LC | |
| <i>Pica pica</i> | 13 | 10.24 | 0.276 | 1.152 | 0.012 | II B | Non-SPEC | LC | |
| <i>Pyrrhonorax pyrrhonorax</i> | 13 | 10.24 | 0.512 | 2.836 | 0.022 | I | SPEC3 | NT | VU |
| <i>Corvus monedula</i> | 3 | 2.36 | 0.031 | 0.216 | 0.001 | II B | Non-SPEC ^E | LC | |
| <i>Corvus cornix</i> | 58 | 45.67 | 1.252 | 2.743 | 0.054 | | Non-SPEC | LC | |
| <i>Corvus corax</i> | 9 | 7.09 | 0.126 | 0.577 | 0.005 | | Non-SPEC | LC | EN |
| <i>Sturnus vulgaris</i> | 11 | 8.66 | 0.913 | 5.947 | 0.039 | II B | SPEC3 | LC | |
| <i>Passer italiae</i> | 11 | 8.66 | 0.717 | 2.806 | 0.031 | | SPEC3 | VU | |
| <i>Passer montanus</i> | 7 | 5.51 | 0.268 | 1.950 | 0.012 | | SPEC3 | VU | |
| <i>Petronia petronia</i> | 3 | 2.36 | 0.079 | 0.586 | 0.003 | | Non-SPEC | LC | VU |
| <i>Montifringilla nivalis</i> * | - | - | - | - | - | | Non-SPEC | LC | EN |
| <i>Fringilla coelebs</i> | 84 | 66.14 | 1.142 | 1.207 | 0.049 | | Non-SPEC ^E | LC | |
| <i>Serinus serinus</i> | 12 | 9.45 | 0.118 | 0.391 | 0.005 | | Non-SPEC ^E | LC | |
| <i>Carduelis chloris</i> | 9 | 7.09 | 0.142 | 0.614 | 0.006 | | Non-SPEC ^E | NT | |
| <i>Carduelis carduelis</i> | 9 | 7.09 | 0.102 | 0.415 | 0.004 | | Non-SPEC | NT | |
| <i>Carduelis cannabina</i> | 31 | 24.41 | 0.480 | 1.075 | 0.021 | | SPEC2 | NT | |
| <i>Pyrrhula pyrrhula</i> | 22 | 17.32 | 0.197 | 0.455 | 0.008 | | Non-SPEC | VU | |
| <i>Coccothraustes coccothraustes</i> | 1 | 0.79 | 0.008 | 0.089 | 0 | | Non-SPEC | LC | EN |
| <i>Emberiza citrinella</i> | 4 | 3.15 | 0.031 | 0.175 | 0.001 | | Non-SPEC ^E | LC | |
| <i>Emberiza cirulus</i> | 26 | 20.47 | 0.291 | 0.656 | 0.013 | | Non-SPEC ^E | LC | |
| <i>Emberiza cia</i> | 2 | 1.57 | 0.016 | 0.125 | 0.001 | | SPEC3 | LC | |
| <i>Emberiza calandra</i> | 9 | 7.09 | 0.189 | 0.764 | 0.008 | | SPEC2 | LC | |

Table 2. Breeding bird community parameters: S = Richness, Smean = mean Richness per point count, A = Abundance, H = Diversity, J = Evenness, non Pass = % non-Passeriformes, N dom. = number of dominant species (see Methods).

| | Point counts | S | Smean | A | H | J | non Pass. | N dom. |
|-------------------------|--------------|----|-------|------|------|------|-----------|--------|
| Montagne della Duchessa | 127 | 79 | 11.99 | 23.2 | 3.77 | 0.86 | 24.1 | 3 |

ing order. The subdominant species were: *Phoenicurus ochruros* (other species characteristic of this habitat), *Anthus trivialis*, *Erithacus rubecula* and *Phylloscopus collybita*. The good abundance of the last three species suggests the influence of ecotonal areas and/or the presence of arboreal elements in this habitat.

The values of bird community parameters of the four habitat types highlight that in the arable crops Richness of species, mean Richness of species per point count, Abundance of individuals and Diversity index reach the highest values (Tab. 4). However, in this habitat the number of dominant species is minimum. The bushes and alpine meadows show the lowest values for all considered parameters. As far as forest environments are concerned, beech wood presents more species than oak wood, but less individuals and a lower value of non-Passeriformes percentage (Tab. 4).

Statistical analysis confirmed the majority of these results in particular the differences between alpine meadows/bushes and the other three habitats. The mean richness of species per point count was significantly different among the four habitats ($H_{3,127} = 52.3$, $P < 0.00001$). In particular the richness resulted significantly lower in the alpine meadows/bushes as compared to oak wood ($Z = 5.48$, $P < 0.00001$), beech wood ($Z = 5.33$, $P < 0.00001$) and arable crops ($Z = 6.11$, $P < 0.00001$). Other pairwise comparisons for this parameter were not significant. Abundance resulted significantly different among the four habitats ($H_{3,127} = 59.2$, $P < 0.00001$) in particular it was significantly lower in alpine meadows/bushes than in oak wood ($Z = 5.73$, $P < 0.00001$), beech wood ($Z = 4.40$, $P < 0.00006$) and arable crops ($Z = 6.92$, $P < 0.00001$); in addition it resulted lower in beech wood as compared to arable crops ($Z = 3.40$, $P < 0.004$). Other pairwise comparisons for this parameter were not significant.

The Dominance values were significantly different among the four habitats ($\chi^2_{3,73} = 8.96$, $P < 0.03$), as well as between arable crops and alpine meadows/bushes ($T = 675$, $P < 0.035$). Other pairwise comparisons for this parameter were not significant.

Non-Passeriformes percentage did not result significantly different among the four habitats and in pairwise comparisons.

DISCUSSION

Results highlight that Montagne della Duchessa Regional Nature Reserve and its surrounding areas present a breeding bird community characterized by values of diversity and richness of species among the highest recorded in Apennine regional areas (De Felici & Sorace 2011). This is likely due to the presence of large sectors of the reserve that show high nature conservation status and to the high level of habitat heterogeneity. However, it has to be noticed that local bird community is also clearly affected by environmental changes produced in the past by man; indeed generalist species, which are successful to settle down in environments modified by anthropogenic activities, are dominant and the most widespread in large parts of study area.

Data collected on environmental preferences of the species and, among them, those dominant in the four main habitat types of the study area, agree with the available information in literature for them (Brunelli et al. 2011) and with investigations carried out in other montane Apennine areas. In particular, as far as alpine meadows are concerned, it is interesting that the six species dominant in the breeding bird community of Campo Imperatore (Gran Sasso d'Italia, 33 km far away from the study area) included *Pyrrhocorax pyrrhocorax* and the same five species which were dominant in the present study (Strinella et al. 2011; for the Simbruini Mts. area, see Vuerich et al. 2006). The highest number of species observed in the arable crops has to be likely related to three factors: the availability of environmental enrichments (trees, bushes, buildings) inside this habitat; the patches have reduced surface leading to detect species living in the adjacent woods and bushes; arable crops are placed at the lowest altitude of the Reserve. The recording of a high richness of species in open habitats due to "edge effect" was also reported in other studies (Farina & Martelli 1979, Sorace 1996, Giraud 2001, Farina 2003, Lorenzetti et al. 2004, Sorace 2011, Bernoni et al. 2012). Species richness and abundance of individuals tend to decrease with altitude (Begon et al. 1986, Sarrocco & Sorace 1997).

Without environmental enrichments and edge effect, breeding bird community of open habitats (meadows, pas-

Breeding bird community of the Montagne della Duchessa

Table 3. Mean number of individuals per point count (\pm SD) and dominance values in each habitat type. Dominant ($\pi > 0.05$) and sub-dominant species ($\pi > 0.02$) in bold and italic, respectively.

| Species | Arable crops | | | Oak wood | | | Beech wood | | | Bushes and meadows | | |
|--|--------------|-------|--------------|----------|-------|--------------|------------|-------|--------------|--------------------|-------|--------------|
| | Mean | SD | π | Mean | SD | π | Mean | SD | π | Mean | SD | π |
| <i>Coturnix coturnix</i> | 0.087 | 0.288 | 0.004 | - | - | - | - | - | - | - | - | - |
| <i>Pernis apivorus</i> | - | - | - | 0.045 | 0.213 | 0.003 | - | - | - | - | - | - |
| <i>Gyps fulvus</i> | - | - | - | - | - | - | - | - | - | 0.028 | 0.167 | 0.005 |
| <i>Accipiter nisus</i> | 0.043 | 0.209 | 0.002 | - | - | - | 0.022 | 0.147 | 0.002 | - | - | - |
| <i>Buteo buteo</i> | - | - | - | 0.091 | 0.294 | 0.006 | 0.109 | 0.482 | 0.010 | - | - | - |
| <i>Falco tinnunculus</i> | 0.087 | 0.288 | 0.004 | - | - | - | - | - | - | 0.028 | 0.167 | 0.005 |
| <i>Columba livia</i> f. <i>domestica</i> | 0.087 | 0.288 | 0.004 | - | - | - | - | - | - | - | - | - |
| <i>Columba palumbus</i> | 0.304 | 0.765 | 0.013 | 0.273 | 0.456 | 0.019 | 0.217 | 0.554 | 0.020 | 0.028 | 0.167 | 0.005 |
| <i>Streptopelia decaocto</i> | 0.087 | 0.417 | 0.004 | - | - | - | - | - | - | - | - | - |
| <i>Streptopelia turtur</i> | - | - | - | 0.045 | 0.213 | 0.003 | - | - | - | - | - | - |
| <i>Cuculus canorus</i> | - | - | - | 0.136 | 0.468 | 0.010 | 0.043 | 0.206 | 0.004 | - | - | - |
| <i>Apus apus</i> | 0.043 | 0.209 | 0.002 | - | - | - | - | - | - | - | - | - |
| <i>Jynx torquilla</i> | 0.087 | 0.288 | 0.004 | - | - | - | - | - | - | - | - | - |
| <i>Picus viridis</i> | 0.087 | 0.288 | 0.004 | 0.182 | 0.395 | 0.013 | 0.065 | 0.250 | 0.006 | - | - | - |
| <i>Dendrocopos major</i> | 0.130 | 0.344 | 0.006 | 0.136 | 0.351 | 0.010 | 0.065 | 0.327 | 0.006 | - | - | - |
| <i>Dendrocopos minor</i> | 0.043 | 0.209 | 0.002 | - | - | - | 0.022 | 0.147 | 0.002 | - | - | - |
| <i>Galerida cristata</i> | 0.478 | 0.947 | <i>0.021</i> | - | - | - | - | - | - | - | - | - |
| <i>Lullula arborea</i> | 0.087 | 0.417 | 0.004 | - | - | - | - | - | - | 0.028 | 0.167 | 0.005 |
| <i>Alauda arvensis</i> | 0.174 | 0.491 | 0.008 | - | - | - | - | - | - | 0.861 | 1.073 | 0.142 |
| <i>Hirundo rustica</i> | 0.304 | 1.105 | 0.013 | - | - | - | - | - | - | - | - | - |
| <i>Delichon urbicum</i> | 0.174 | 0.650 | 0.008 | 0.045 | 0.213 | 0.003 | - | - | - | - | - | - |
| <i>Anthus campestris</i> | - | - | - | - | - | - | - | - | - | 0.417 | 0.649 | 0.068 |
| <i>Anthus trivialis</i> | - | - | - | 0.045 | 0.213 | 0.003 | 0.109 | 0.315 | 0.010 | 0.167 | 0.447 | <i>0.027</i> |
| <i>Anthus spinoletta</i> | - | - | - | - | - | - | - | - | - | 0.889 | 0.854 | 0.146 |
| <i>Motacilla alba</i> | 0.174 | 0.388 | 0.008 | - | - | - | - | - | - | 0.056 | 0.232 | 0.009 |
| <i>Troglodytes troglodytes</i> | 0.130 | 0.458 | 0.006 | 0.227 | 0.429 | 0.016 | 0.543 | 0.657 | <i>0.049</i> | 0.083 | 0.28 | 0.014 |
| <i>Erithacus rubecula</i> | 1.000 | 1.206 | <i>0.043</i> | 1.955 | 0.950 | 0.139 | 1.630 | 0.853 | 0.146 | 0.139 | 0.487 | <i>0.023</i> |
| <i>Luscinia megarhynchos</i> | 0.478 | 0.593 | <i>0.021</i> | 0.091 | 0.294 | 0.006 | - | - | - | 0.083 | 0.368 | 0.014 |
| <i>Phoenicurus ochruros</i> | 0.087 | 0.288 | 0.004 | - | - | - | 0.022 | 0.147 | 0.002 | 0.194 | 0.467 | <i>0.032</i> |
| <i>Phoenicurus phoenicurus</i> | 0.348 | 0.647 | 0.015 | 0.045 | 0.213 | 0.003 | 0.065 | 0.327 | 0.006 | - | - | - |
| <i>Saxicola torquatus</i> | 0.087 | 0.288 | 0.004 | - | - | - | - | - | - | 0.083 | 0.368 | 0.014 |
| <i>Oenanthe oenanthe</i> | - | - | - | - | - | - | - | - | - | 1.167 | 1.028 | 0.192 |
| <i>Monticola saxatilis</i> | - | - | - | - | - | - | - | - | - | 0.083 | 0.368 | 0.014 |
| <i>Turdus merula</i> | 0.565 | 0.788 | <i>0.025</i> | 1.364 | 1.590 | 0.097 | 0.457 | 0.504 | <i>0.041</i> | 0.028 | 0.167 | 0.005 |
| <i>Turdus philomelos</i> | 0.043 | 0.209 | 0.002 | - | - | - | 0.022 | 0.147 | 0.002 | - | - | - |
| <i>Turdus viscivorus</i> | - | - | - | 0.045 | 0.213 | 0.003 | 0.239 | 0.480 | <i>0.021</i> | - | - | - |
| <i>Sylvia atricapilla</i> | 1.087 | 1.240 | <i>0.047</i> | 1.182 | 1.006 | 0.084 | 0.978 | 0.802 | 0.088 | 0.083 | 0.280 | 0.014 |
| <i>Sylvia cantillans</i> | 0.130 | 0.458 | 0.006 | - | - | - | - | - | - | 0.056 | 0.232 | 0.009 |
| <i>Phylloscopus bonelli</i> | 0.087 | 0.288 | 0.004 | 0.500 | 0.802 | <i>0.035</i> | 0.283 | 0.544 | <i>0.025</i> | - | - | - |
| <i>Phylloscopus sibilatrix</i> | - | - | - | - | - | - | 0.130 | 0.542 | 0.012 | - | - | - |
| <i>Phylloscopus collybita</i> | 0.261 | 0.541 | 0.011 | 0.909 | 0.868 | 0.065 | 0.674 | 0.790 | 0.061 | 0.167 | 0.447 | <i>0.027</i> |
| <i>Regulus ignicapilla</i> | - | - | - | 0.182 | 0.501 | 0.013 | 0.065 | 0.250 | 0.006 | - | - | - |
| <i>Ficedula albicollis</i> | - | - | - | - | - | - | 0.043 | 0.206 | 0.004 | - | - | - |
| <i>Aegithalos caudatus</i> | 0.087 | 0.288 | 0.004 | 0.318 | 0.568 | <i>0.023</i> | 0.109 | 0.434 | 0.010 | - | - | - |
| <i>Cyanistes caeruleus</i> | 0.739 | 0.964 | <i>0.032</i> | 1.273 | 1.032 | 0.090 | 0.717 | 0.750 | 0.064 | 0.111 | 0.398 | 0.018 |
| <i>Parus major</i> | 0.913 | 1.083 | <i>0.040</i> | 1.091 | 1.192 | 0.077 | 0.522 | 0.752 | <i>0.047</i> | 0.028 | 0.167 | 0.005 |

continued

| Species | Arable crops | | | Oak wood | | | Beech wood | | | Bushes and meadows | | |
|--------------------------------|--------------|-------|--------------|----------|-------|--------------|------------|-------|--------------|--------------------|-------|--------------|
| | Mean | SD | pi | Mean | SD | pi | Mean | SD | pi | Mean | SD | pi |
| <i>Periparus ater</i> | 0.217 | 0.736 | 0.009 | 0.136 | 0.468 | 0.010 | 0.543 | 1.048 | 0.049 | 0.028 | 0.167 | 0.005 |
| <i>Poecile palustris</i> | 0.261 | 0.689 | 0.011 | 0.864 | 1.167 | 0.061 | 0.804 | 0.885 | 0.072 | - | - | - |
| <i>Sitta europaea</i> | 0.304 | 0.635 | 0.013 | 0.682 | 1.129 | 0.048 | 0.717 | 0.911 | 0.064 | - | - | - |
| <i>Certhia familiaris</i> | - | - | - | - | - | - | 0.087 | 0.285 | 0.008 | - | - | - |
| <i>Certhia brachydactyla</i> | 0.087 | 0.288 | 0.004 | 0.045 | 0.213 | 0.003 | 0.065 | 0.327 | 0.006 | - | - | - |
| <i>Oriolus oriolus</i> | 0.043 | 0.209 | 0.002 | 0.045 | 0.213 | 0.003 | - | - | - | - | - | - |
| <i>Lanius collurio</i> | 0.174 | 0.491 | 0.008 | - | - | - | - | - | - | 0.028 | 0.167 | 0.005 |
| <i>Garrulus glandarius</i> | 0.478 | 0.994 | 0.021 | 0.636 | 0.902 | 0.045 | 0.261 | 0.612 | 0.023 | 0.028 | 0.167 | 0.005 |
| <i>Pica pica</i> | 0.696 | 1.295 | 0.030 | - | - | - | - | - | - | - | - | - |
| <i>Pyrrhocorax pyrrhocorax</i> | - | - | - | - | - | - | - | - | - | 0.056 | 0.232 | 0.009 |
| <i>Corvus monedula</i> | 0.130 | 0.458 | 0.006 | - | - | - | - | - | - | - | - | - |
| <i>Corvus cornix</i> | 1.478 | 2.391 | 0.064 | 0.227 | 0.685 | 0.016 | 0.022 | 0.147 | 0.002 | 0.056 | 0.232 | 0.009 |
| <i>Corvus corax</i> | - | - | - | - | - | - | - | - | - | 0.111 | 0.523 | 0.018 |
| <i>Sturnus vulgaris</i> | 3.087 | 8.404 | 0.134 | - | - | - | - | - | - | - | - | - |
| <i>Passer italiae</i> | 3.391 | 5.168 | 0.147 | - | - | - | - | - | - | - | - | - |
| <i>Passer montanus</i> | 1.435 | 4.470 | 0.062 | - | - | - | - | - | - | - | - | - |
| <i>Petronia petronia</i> | 0.174 | 0.576 | 0.008 | - | - | - | - | - | - | - | - | - |
| <i>Fringilla coelebs</i> | 0.304 | 0.559 | 0.013 | 0.727 | 0.935 | 0.052 | 1.022 | 1.064 | 0.092 | 0.111 | 0.398 | 0.018 |
| <i>Serinus serinus</i> | 0.304 | 0.635 | 0.013 | - | - | - | - | - | - | 0.056 | 0.232 | 0.009 |
| <i>Carduelis chloris</i> | 0.304 | 0.876 | 0.013 | - | - | - | 0.043 | 0.295 | 0.004 | - | - | - |
| <i>Carduelis carduelis</i> | 0.348 | 0.714 | 0.015 | 0.091 | 0.294 | 0.006 | - | - | - | 0.028 | 0.167 | 0.005 |
| <i>Carduelis cannabina</i> | 0.348 | 0.935 | 0.015 | 0.136 | 0.351 | 0.010 | - | - | - | 0.722 | 1.446 | 0.119 |
| <i>Pyrrhula pyrrhula</i> | - | - | - | 0.182 | 0.395 | 0.013 | 0.304 | 0.511 | 0.027 | - | - | - |
| <i>Emberiza citrinella</i> | 0.043 | 0.209 | 0.002 | - | - | - | 0.043 | 0.206 | 0.004 | - | - | - |
| <i>Emberiza cirrus</i> | 0.304 | 0.559 | 0.013 | 0.136 | 0.351 | 0.010 | 0.022 | 0.147 | 0.002 | 0.056 | 0.232 | 0.009 |
| <i>Emberiza cia</i> | - | - | - | - | - | - | 0.043 | 0.206 | 0.004 | - | - | - |
| <i>Emberiza calandra</i> | 0.522 | 0.846 | 0.023 | - | - | - | - | - | - | - | - | - |

tures, bushes) are usually characterized by a low number of species and individuals, with few dominant species yielding to low values of Diversity and Evenness indices (Farina 1982, Sorace 1996; for the negative relation between degree of vegetation cover and Evenness, see for example Symonds & Johnson 2008). Conversely, non-Passeriformes percentage usually increases in the ecological succession toward climax community (Ferry & Frochot 1970).

Regarding species of conservation interest, the Montagne della Duchessa Reserve may play a role of great importance for *Alectoris graeca* and *Gyps fulvus*. In this area, the former species achieves density values among the highest of Apennines (Sorace et al. 2011, 2013); the latter is present with a colony of 10 breeding pairs, that established following a reintroduction project started in 1994 (Allavena & Panella 2000, Allavena 2012).

In spite of its reduced surface, the study area has a great value at regional scale, since some of the most local-

ized species are breeding inside it. In fact, besides the two above quoted species, *Tichodroma muraria*, *Pyrrhocorax pyrrhocorax*, *Petronia petronia*, *Montifringilla nivalis* and *Coccothraustes coccothraustes* are found (Brunelli et al. 2011).

In order to protect local species of conservation concern (see Tab. 1) in a more effective manner, an enlargement of Montagne della Duchessa Regional Nature Reserve would be appropriate. In particular the inclusion of the whole massif of Cava Mountain should benefit, for example, species such as *Alectoris graeca* and *Anthus campestris* as well as *Aquila chrysaetos* that nests in areas very proximate to the study area and regularly uses alpine meadows of the massif for feeding activities. Moreover, the insertion of Corvaro plain should favour species such as *Lullula arborea*, *Anthus campestris* and *Lanius collurio* as well as *Pyrrhocorax pyrrhocorax*, that feeds in this area during the winter period.

Table 4. Breeding bird community parameters for each habitat type: S = Richness, Smean = Mean Richness per point count, A = Abundance, H = Diversity, J = Evenness, non Pass. = % non-Passeriformes; N dom. = number of dominant species (see Methods).

| Habitat | Point counts | S | Smean | A | H | J | non Pass. | N dom. |
|--------------------|--------------|----|-------|-------|------|------|-----------|--------|
| Arable crops | 23 | 54 | 10.65 | 22.96 | 3.31 | 0.83 | 20.37 | 4 |
| Oak wood | 22 | 34 | 9.36 | 14.09 | 2.96 | 0.84 | 20.59 | 8 |
| Beech wood | 46 | 37 | 8.20 | 11.13 | 2.99 | 0.83 | 18.92 | 7 |
| Bushes and Meadows | 36 | 33 | 4.22 | 6.08 | 2.72 | 0.78 | 9.10 | 5 |

Acknowledgements – The authors would like to thank the Montagne della Duchessa Regional Nature Reserve for funding the project. We thank for their help and support Dr. Gianluca Scialanga, Director of the Montagne della Duchessa Regional Nature Reserve, Dr. Silvia Scozzafava and the staff of the Reserve, Dr. Emanuela Peria of the Regional Park Agency of the Latium Region.

REFERENCES

- AA.VV., 2004. Piano di Gestione dei proposti Siti di Interesse Comunitario IT6020020 Monti della Duchessa (area sommitale), IT6020021 Monte Duchessa (Vallone Cieco e Bosco Cartore), e della Zona di Protezione Speciale IT6020046 Riserva Naturale “Montagne della Duchessa”. Documento tecnico non pubblicato.
- Allavena S., 2012. Grifone *Gyps fulvus*. In: Aradis A., Sarrocco S. & Brunelli M. Analisi dello status e della distribuzione dei rapaci diurni nidificanti nel Lazio. ISPRA. Quaderni Natura e BioDiversity 2/2012: 36-41.
- Allavena S. & Panella M., 2000. La reintroduzione del Grifone *Gyps fulvus* nella Riserva Naturale del Monte Velino. Alula 7: 10-19.
- Allavena S., Panella M. & Spinetti M., 1999. La reintroduzione del Corvo imperiale *Corvus c. corax* nella Riserva Naturale Orientata del Monte Velino-Montagna della Duchessa. Alula 6: 131-136.
- Apollonio M., Baldaccini N.E., Blasi C., Bulgarini F., Celada., Cerfolli F., Cerra M., Corona P.M., Dell’Anna L., Feoli E., Gatto M., Guifa C., La Posta A., Maggiore A.M., Manes F., Marchetti M., Morabito A., Paris G., Renzi F., Rizzi V., Rossi R., Rubino T., Solinas M. & Tartaglini N., 2004. La gestione dei siti Natura 2000 in Italia: linee guida e orientamenti gestionali. Quaderni Conservazione Natura 18: 213-224.
- Battisti C., Bologna M.A. & Vignoli L., 2010. Breeding birds in an Apennine massif (Majella, central Italy): do “common species” could act as surrogate for characterize species richness and composition of the communities? *Ecológia* (Bratislava) 29: 207-218.
- Battisti C. & Guidi A., 2012. Gli uccelli nidificanti nella Riserva Naturale di Monte Catillo. Check-list, distribuzione locale e status di conservazione. Provincia di Roma, Assessorato alle Politiche dell’Agricoltura, 144 pp.
- Belmaker J., Sekercioglu C.H. & Jetz W., 2012. Global patterns of specialization and coexistence in bird assemblages. *Journal Biogeography* 39: 193-203.
- Begon M., Harper J.L. & Townsend C.R., 1986. Ecology, Individuals, Populations and Communities. Blackwell Scientific Publications, London, 876 pp.
- Bernoni M., Brunelli M., De Santis E., Sarrocco S. & Cecere J.G., 2009. Distribuzione e stima delle popolazioni di Gracchio corallino, *Pyrhcorax pyrrhcorax*, e di Gracchio alpino *Pyrhcorax graculus*, nel Lazio. *Rivista italiana Ornitologia* 79: 11-21.
- Bernoni M., Sorace A., Cecere J.C., Biondi M., Cento M., Guerrieri G. & Savo E., 2012. La comunità degli uccelli nidificanti nella ZPS IT-603005 “Comprensorio Tolfetano-Cerite-Manziate”. *Alula* 19: 11-27.
- Bibby C.J., Burgess N.D., Hill D.A. & Mustoe S.H., 2000. Bird Census Techniques. II ed. Academic Press, London.
- BirdLife International, 2004. Birds in Europe: population estimates, trends and conservation status. BirdLife International Conservation Series 12, Cambridge, UK.
- Blondel J., Ferry C. & Frochet B., 1970. La méthode des indices ponctuels d’abondance (IPA) ou des relevés d’avifaune par “stations d’écoute”. *Alauda* 38: 55-71.
- Brandmayr P., Cagnin M., Mingozzi T. & Pizzolotto R., 1996. Map of Zoocoenoses and Evaluation for the Menta River Dam in Aspromonte (Calabria, Italy). *Z. Ökologie u. Naturschutz*, 5: 15-28.
- Brunelli M., Sarrocco S., Corbi F., Sorace A., Boano A., De Felici S., Guerrieri G., Meschini A. & Roma S. (a cura di), 2011. Nuovo Atlante degli uccelli nidificanti nel Lazio. Edizioni ARP (Agenzia Regionale Parchi), Roma, 464 pp.
- Calvario E., Brunelli M., Sarrocco S., Bulgarini F., Fraticelli F. & Sorace A., 2011. Lista Rossa degli Uccelli nidificanti nel Lazio (2010). In: Brunelli M., Sarrocco S., Corbi F., Sorace A., Boano A., De Felici S., Guerrieri G., Meschini A. & Roma S. (a cura di), Nuovo Atlante degli uccelli nidificanti nel Lazio. Edizioni ARP (Agenzia Regionale Parchi), Roma: 427-435.
- D.R.E.A.M. Italia, 2003. Piano di assestamento forestale della Riserva Naturale Regionale Montagne della Duchessa. Documento tecnico non pubblicato.
- De Felici S. & Sorace A., 2011. Analisi della ricchezza di specie. In: Brunelli M., Sarrocco S., Corbi F., Sorace A., Boano A., De Felici S., Guerrieri G., Meschini A. & Roma S. (a cura di), Nuovo Atlante degli uccelli nidificanti nel Lazio. Edizioni ARP (Agenzia Regionale Parchi), Roma: 409-416.
- De Filippo G., Caputo V. & Kalby M., 1985. Le comunità di uccelli in una fustaia di faggio sui Monti Alburni (Sud Italia). *Bollettino Società Naturalisti Napoli* 94: 221-227.
- De Filippo G. & Kalby M., 1985. Modificazioni nella struttura di una comunità di uccelli in seguito allo sfruttamento silvo-culturale di una fustaia di faggio. In: Fasola M. (red.). *Atti III Convegno italiano Ornitologia*, Pavia: 198-200.
- De Santis E., Sorace A. & Tarquini L., 2009. L’Atlante degli uccelli nidificanti nel Parco Naturale Regionale dei Monti Simbruini (Lazio, Italia centrale): dati preliminari. *Alula* 16: 541-543.
- Devictor V. & Robert A., 2009. Measuring community responses to large-scale disturbance in conservation biogeography. *Diversity Distribution* 15: 122-130.
- Di Carlo E.A., 1980. Indagine preliminare sulla presenza passata ed attuale dell’Aquila reale, *Aquila chrysaetos*, sugli Appennini. *Gli Uccelli d’Italia* 5: 263-283.

- Farina A., 1982. Bird community of Mediterranean Forest of Migliarino (Pisa - central Italy). *Avocetta* 6: 75-81.
- Farina A., 2003. *Ecologia del paesaggio. Principi, metodi e applicazioni*. UTET Libreria, Torino, 674 pp.
- Farina A. & Martelli C., 1979. Breeding bird census of an Italian mediterranean habitat: the Parco naturale della Maremma. VI International Congress Bird Census Work: 129-135.
- Ferry C. & Frochot B. 1970. L'avifaune nidificatrice d'une forêt de chênes pedunculés en Bourgogne: étude de deux successions écologiques. *Terre et Vie* 2: 153-250.
- Fornasari L., Londi G., Buvoli L., Tellini Florenzano G., La Gioia G., Pedrini P., Bricchetti P. & De Carli E., 2010. Distribuzione geografica e ambientale degli uccelli comuni in Italia, 2000-2004 (dati del progetto MITO2000). *Avocetta* 34: 5-224.
- Fulco E. & Tellini Florenzano G., 2008. Composizione e struttura della comunità ornitica nidificante in una faggeta della Basilicata. *Avocetta* 32: 55-60.
- Giraud L., 2001. Relazione fra fattori ambientali, stagionali e avifauna del piano montano. *Avocetta* 25: 54.
- Lloyd M. & Ghelardi R., 1964. A table for calculating the "Equitability" component of species diversity. *Journal Animal Ecology* 33: 217-225.
- Lorenzetti E., Ukmar E. & Battisti C., 2004. Comunità ornitiche nidificanti nel Sito di Importanza Comunitaria "Monte Guadagnolo" (Monti Prenestini, Italia centrale). *Alula* 11: 105-112.
- Mancinelli A., De Castro G., De Sanctis A. & Posillico M., 2009. La comunità ornitica delle praterie d'altitudine di Monte Greco, Appennino centrale. *Alula* 16: 366-368.
- Mingozzi T., 1994. L'Avifauna della Riserva Naturale Orientata "Valle del fiume Argentino" (Orsomarso, CS): Analisi faunistico-ecologica ed indicazioni conservazionistico-gestionali. Collana Verde, Ministero Risorse Agricole, Alimentari e Forestali, Roma, 96: 1-94.
- Papi R., Luiselli L. & Rugiero L., 2010. Atlante dell'avifauna e dell'erpeto-fauna della Riserva Naturale Regionale Monte Rufeno. Collana Atlanti Locali, Edizioni ARP, Roma.
- Peronace V., Cecere J.C., Gustin M. & Rondinini C., 2012. Lista Rossa 2011 degli Uccelli nidificanti in Italia. *Avocetta* 36: 11-58.
- Pielou E.C., 1966. The measurement of diversity in different types of biological collections. *Journal Theoretical Biology* 13: 131-144.
- Primack R.B. & Carotenuto L., 2003. *La conservazione della natura*. Zanichelli, Bologna.
- Regione Lazio, 2010. Carta delle formazioni naturali e seminaturali della Regione Lazio 1:10.000: approfondimento e aggiornamento della Carta dell'Uso del Suolo (CUS Lazio DGR n. 953 del 28 marzo 2000), Agenzia Regionale Parchi - Regione Lazio.
- Quatrini A., Scarfò F. & Zapparoli M., 2012. Atlante degli uccelli nidificanti nella Riserva Naturale Regionale del Lago di Vico (Lazio, Viterbo). *Alula* 19: 97-163.
- Rossi D., 1947. Risultati di ricerche ornitologiche sulle montagne d'Abruzzo. Parte 1. Gruppo Monti Sirente-Velino - Anni 1945-1947. *Rivista italiana Ornitologia* 17: 87-128.
- Russo D. & Cistrone L. & Carotenuto L., 2010. I chiroterteri della Riserva Naturale Regionale Montagne della Duchessa. Collana Atlanti Locali, Edizioni ARP, Roma.
- Sarrocchio S. & Sorace A., 1997. Le comunità ornitiche nidificanti nella Riserva regionale "Lago di Vico". *Rivista italiana Ornitologia* 67: 71-74.
- Shannon C.E. & Weaver W., 1963. *Mathematical theory of communication*. University of Illinois Press, Urbana.
- Sorace A., 1996. Alcuni dati sulle comunità ornitiche del comprensorio Monte Cairo Valle del Melfa (FR). *Alula* 3:17-22.
- Sorace A., 2011. Le comunità ornitiche degli ambienti aperti della Sila Grande (Calabria) con note sull'apparente contrazione demografica di *Saxicola rubetra*, *Regulus regulus* e *Carduelis spinus*. *Avocetta* 35: 53-60.
- Sorace A., Artese C., Antonucci M., Bernoni M., Bonanni M., Brusaferrò A., Carafa M., Carotenuto L., Cortone P., De Filippo G., De Santis E., Forcaoni P., Fabrizio M., Fulco E., Guglielmi S., Latini R., Liberatoscioli E., Magrini M., Mangiacotti M., Mariani F., Pellegrini M., Peria E., Pinchiurri V., Properzi S., Riga F., Scalisi M., Spera M. & Trocchi V., 2013. Status and distribution of Rock Partridge *Alectoris graeca* in Apennine areas. *Avocetta* 37: 111-118.
- Sorace A. & Gustin M., 2010. Bird species of conservation concern along urban gradients in Italy. *Biodiversity and Conservation* 19: 205-221.
- Sorace A., Properzi S., Guglielmi S., Riga F., Trocchi V. & Scalisi M., 2011. La Coturnice nel Lazio: status e piano d'azione. Edizioni ARP, Roma.
- Spinetti M., 1997. Fauna del Parco Regionale Sirente-Velino. Parconatura 2. Gruppo Tipografico Editoriale, L'Aquila.
- Strinella E., Sorace A., Lalli G., Ferrossi D. & Savo E., 2011. Analisi preliminare dei passeriformi nidificanti in praterie d'altitudine appenniniche: l'area di Campo Imperatore (AQ). *Alula* 18: 101-104.
- Sutherland W.J., Newton I. & Green R.E., 2004. *Ecological Census Techniques*. Oxford University Press, Oxford.
- Symonds M. R. E. & Johnson C.N., 2008. Species Richness and Evenness in Australian Birds. *American Naturalist* 171: 480-490.
- Taffon D., Giucca F., Battisti C. (a cura di), 2008. Atlante degli uccelli nidificanti nel Parco Regionale dell'Appia Antica. Ente Parco Regionale dell'Appia Antica - Provincia di Roma, Assessorato alle Politiche dell'Agricoltura e dell'Ambiente, Gangemi Editore, 176 pp.
- Turcek F.J., 1956. Zur Frage der Dominanze in Vogelpopulationen. *Waldhygiene* 8: 249-257.
- Ukmar E., Badaloni D., Taffon D., Lorenzetti E. & Sorace A., 2009. L'Atlante degli uccelli nidificanti nel Parco Regionale dei Castelli Romani. *Alula* 16: 689-691.
- Vuerich V., Bologna M.A. & Battisti C., 2006. Comunità ornitiche nidificanti in tre tipologie vegetazionali lungo un gradiente altitudinale nei Monti Simbruini (Lazio, Appennini centrali) (Aves). *Aldrovandia* 2: 49-53.

Associate editor: **Bruno Massa**